



## The Maximum Number of Edges in a Three-Dimensional Grid-Drawing

*Prosenjit Bose*

School of Computer Science  
Carleton University  
Ottawa, Ontario, Canada  
[jit@scs.carleton.ca](mailto:jit@scs.carleton.ca)

*Jurek Czyzowicz*

Département d'informatique et d'ingénierie  
Université du Québec en Outaouais  
Gatineau, Québec, Canada  
[jurek@uqo.ca](mailto:jurek@uqo.ca)

*Pat Morin*     *David R. Wood*

School of Computer Science  
Carleton University  
Ottawa, Ontario, Canada  
[{morin,davidw}@scs.carleton.ca](mailto:{morin,davidw}@scs.carleton.ca)

### Abstract

An exact formula is given for the maximum number of edges in a graph that admits a three-dimensional grid-drawing contained in a given bounding box. The first universal lower bound on the volume of three-dimensional grid-drawings is obtained as a corollary. Our results generalise to the setting of multi-dimensional polyline grid-drawings.

Article Type	Communicated by	Submitted	Revised
concise paper	G. Liotta	April 2003	March 2004

A *three-dimensional (straight-line) grid-drawing* of a graph represents the vertices by distinct points in  $\mathbb{Z}^3$ , and represents each edge by a line-segment between its endpoints that does not intersect any other vertex, and does not intersect any other edge except at the endpoints. A folklore result states that every (simple) graph has a three-dimensional grid-drawing (see [2]). We therefore are interested in grid-drawings with small ‘volume’.

The *bounding box* of a three-dimensional grid-drawing is the axis-aligned box of minimum size that contains the drawing. If the bounding box has side lengths  $X - 1$ ,  $Y - 1$  and  $Z - 1$ , then we speak of an  $X \times Y \times Z$  grid-drawing with *volume*  $X \cdot Y \cdot Z$ . That is, the volume of a 3D drawing is the number of grid-points in the bounding box. (This definition is formulated to ensure that a two-dimensional grid-drawing has positive volume.) Our main contribution is the following extremal result.

**Theorem 1.** *The maximum number of edges in an  $X \times Y \times Z$  grid-drawing is exactly*

$$(2X - 1)(2Y - 1)(2Z - 1) - XYZ .$$

*Proof.* Let  $B$  the bounding box in an  $X \times Y \times Z$  grid-drawing of a graph  $G$  with  $n$  vertices and  $m$  edges. Let  $P = \{(x, y, z) \in B : 2x, 2y, 2z \in \mathbb{Z}\}$ . Observe that  $|P| = (2X - 1)(2Y - 1)(2Z - 1)$ . The midpoint of every edge of  $G$  is in  $P$ , and no two edges share a common midpoint. Hence  $m \leq |P|$ . In addition, the midpoint of an edge does not intersect a vertex. Thus

$$m \leq |P| - n . \tag{1}$$

A drawing with the maximum number of edges has no edge that passes through a grid-point. Otherwise, sub-divide the edge, and place a new vertex at that grid-point. Thus  $n = XYZ$ , and  $m \leq |P| - XYZ$ , as claimed.

This bound is attained by the following construction. Associate a vertex with each grid-point in an  $X \times Y \times Z$  grid-box  $B$ . As illustrated in Figure 1, every vertex  $(x, y, z)$  is adjacent to each of  $(x \pm 1, y, z)$ ,  $(x, y \pm 1, z)$ ,  $(x, y, z \pm 1)$ ,  $(x + 1, y + 1, z)$ ,  $(x - 1, y - 1, z)$ ,  $(x + 1, y, z + 1)$ ,  $(x - 1, y, z - 1)$ ,  $(x, y + 1, z + 1)$ ,  $(x, y - 1, z - 1)$ ,  $(x + 1, y + 1, z + 1)$ , and  $(x - 1, y - 1, z - 1)$ , unless such a grid-point is not in  $B$ . It is easily seen that no two edges intersect, except at a common endpoint. Furthermore, every point in  $P$  is either a vertex or the midpoint of an edge. Thus the number of edges is  $|P| - XYZ$ .  $\square$

Theorem 1 can be interpreted as a lower bound on the volume of a three-dimensional grid-drawing of a given graph. Many upper bounds on the volume of three-dimensional grid-drawings are known [1–6, 8, 10, 11, 14, 15]. There are two known non-trivial lower bounds for specific families of graphs. Cohen, Eades, Lin, and Ruskey [2] proved that the minimum volume of a three-dimensional grid-drawing of the complete graph  $K_n$  is  $\Theta(n^3)$ . The lower bound follows from the fact that  $K_5$  is not planar, and hence at most four vertices can lie in a single grid-plane. The second lower bound is due to Pach, Thiele, and Tóth [14], who proved that the minimum volume of a three-dimensional grid-drawing




Figure 1: The neighbourhood of a vertex.

of the complete bipartite graph  $K_{n,n}$  is  $\Theta(n^2)$ . The proof of the lower bound is based on the observation that no two edges from one colour class to the other have the same direction. The result follows since the number of distinct vectors between adjacent vertices is at most a constant times the volume. (Calamoneri and Sterbini [1] had earlier proved that every three-dimensional grid-drawing of  $K_{n,n}$  has  $\Omega(n^{3/2})$  volume.) The following corollary of Theorem 1 generalises the lower bound of Pach *et al.* [14] to all graphs.

**Corollary 1.** *A three-dimensional grid-drawing of a graph with  $n$  vertices and  $m$  edges has volume greater than  $(m + n)/8$ .*

*Proof.* Let  $v$  be the volume of an  $X \times Y \times Z$  grid-drawing. By (1),  $m \leq |P| - n < 8v - n$ , and hence  $v > (m + n)/8$ .  $\square$

Theorem 1 generalises to multi-dimensional polyline grid-drawings (where edges may bend at grid-points) as follows. Note that upper bounds for the volume of three-dimensional polyline grid-drawings have been recently established [7, 9, 13].

**Theorem 2.** *Let  $B \subseteq \mathbb{R}^d$  be a convex set. Let  $S = B \cap \mathbb{Z}^d$  be the set of grid-points in  $B$ . The maximum number of edges in a polyline grid-drawing with bounding box  $B$  is at most  $(2^d - 1)|S|$ . If  $B$  is an  $X_1 \times \dots \times X_d$  grid-box, then the maximum number of edges is exactly*

$$\prod_{i=1}^d (2X_i - 1) - \prod_{i=1}^d X_i .$$

*Proof.* Let  $P = \{x \in B : 2x \in \mathbb{Z}^d\}$ . Consider a polyline grid-drawing with bounding box  $B$ . The midpoint of every segment is in  $P$ , and no two segments share a common midpoint. A drawing with the maximum number of edges has no edge that passes through a grid-point. Otherwise, sub-divide the edge, and place a new vertex at that grid-point. Thus the number of segments, and hence the number of edges, is at most  $|P| - |S| \leq (2^d - 1)|S|$ .

If  $B$  is an  $X_1 \times \cdots \times X_d$  grid-box, then  $|P| - |S| = \prod_{i=1}^d (2X_i - 1) - \prod_{i=1}^d X_i$ . To construct a grid-drawing with this many edges, associate one vertex with each grid-point in  $S$ . Observe that every point  $x \in P \setminus S$  is in the interior of exactly one unit-sized  $d'$ -dimensional hypercube with corners in  $S$ , where  $1 \leq d' \leq d$ . For every point  $x \in P \setminus S$ , add an edge passing through  $x$  between one pair of opposite vertices of the unit-sized hypercube corresponding to  $x$ . Edges only intersect at common endpoints, since these unit-sized hypercubes only intersect along their boundaries. Every point in  $P$  contains a vertex or a midpoint of an edge. Thus the number of edges is precisely  $|P| - |S|$ .  $\square$

## References

- [1] T. Calamoneri and A. Sterbini. 3D straight-line grid drawing of 4-colorable graphs. *Inform. Process. Lett.*, **63**(2):97–102, 1997.
- [2] R. F. Cohen, P. Eades, T. Lin, and F. Ruskey. Three-dimensional graph drawing. *Algorithmica*, **17**(2):199–208, 1996.
- [3] E. Di Giacomo. Drawing series-parallel graphs on restricted integer 3D grids. In [12], pp. 238–246.
- [4] E. Di Giacomo, G. Liotta, and S. Wismath. Drawing series-parallel graphs on a box. In *Proc. 14th Canadian Conf. on Computational Geometry (CCCG '02)*, pp. 149–153, The University of Lethbridge, Canada, 2002.
- [5] E. Di Giacomo and H. Meijer. Track drawings of graphs with constant queue number. In [12], pp. 214–225.
- [6] V. Dujmović, P. Morin, and D. R. Wood. Layout of graphs with bounded tree-width, *SIAM J. Comput.*, accepted in 2004.
- [7] V. Dujmović and D. R. Wood. Layouts of graph subdivisions, In *Proc. of 12th International Symposium on Graph Drawing (GD '04), Lecture Notes in Comput. Sci.*, Springer, to appear.
- [8] V. Dujmović and D. R. Wood. Three-dimensional grid drawings with sub-quadratic volume. In [12], pp. 190–201. Also in J. Pach, ed., *Towards a Theory of Geometric Graphs*, vol. 342 of *Contemporary Mathematics*, pp. 55–66, Amer. Math. Soc., 2004.
- [9] B. Dyck, J. Joevenazzo, E. Nickle, J. Wilsdon, and S. K. Wismath. Drawing  $K_n$  in three dimensions with two bends per edge. Tech. Rep. TR-CS-01-04, Department of Mathematics and Computer Science, University of Lethbridge, 2004.
- [10] S. FELSNER, G. LIOTTA, AND S. WISMATH, Straight-line drawings on restricted integer grids in two and three dimensions. *J. Graph Algorithms Appl.*, **7**(4):363–398, 2003.
- [11] T. Hasunuma. Laying out iterated line digraphs using queues. In [12], pp. 202–213.
- [12] G. Liotta. ed., *Proc. 11th International Symp. on Graph Drawing (GD '03)*, vol. 2912 of *Lecture Notes in Comput. Sci.*, Springer, 2004.
- [13] P. Morin, and D. R. Wood. Three-Dimensional 1-Bend Graph Drawings. In *Proc. 16th Canadian Conference on Computational Geometry (CCCG '04)*, Concordia University, Canada, 2004.

- [14] J. Pach, T. Thiele, and G. Tóth. Three-dimensional grid drawings of graphs. In G. Di Battista. ed., *Proc. 5th International Symp. on Graph Drawing (GD '97)*, vol. 1353 of *Lecture Notes in Comput. Sci.*, pp. 47–51, Springer, 1997. Also in B. Chazelle. J. E. Goodman, and R. Pollack, eds., *Advances in discrete and computational geometry*, vol. 223 of *Contemporary Mathematics*, pp. 251–255, Amer. Math. Soc., 1999.
- [15] T. Poranen. A new algorithm for drawing series-parallel digraphs in 3D. Tech. Rep. A-2000-16, Dept. of Computer and Information Sciences, University of Tampere, Finland, 2000.